

TOPIC: Staff Resources	SUBJECT: COVID-19 Reference Document for Symptoms Ver. 7	Code: SR000
		Date of Issue: September 21,2020
		Revised:
		Page: Page 1 of 3



Ministry of Health

COVID-19 Reference Document for Symptoms

Version 7.0 – September 21, 2020

This document outlines the symptoms, signs, and clinical features which have been most commonly associated with COVID-19. This information is current as of September 21, 2020 and may be updated as the situation on COVID-19 continues to evolve. If there is a discrepancy between this list and other guidance, this list should be considered as the most up to date.

When assessing for the symptoms below the focus should be on evaluating if they are **new, worsening, or different from an individual’s baseline health status (usual state)**. Symptoms should not be chronic or related to other known causes or conditions (see examples below).

Common symptoms of COVID-19 include:

- **Fever** (temperature of 37.8°C/100.0°F or greater)
- **Cough** (that is new or worsening (e.g. continuous, more than usual if chronic cough) including croup (barking cough, making a whistling noise when breathing)
 - *Not related to other known causes or conditions (e.g., chronic obstructive pulmonary disease)*
- **Shortness of breath** (dyspnea, out of breath, unable to breathe deeply, wheeze, that is worse than usual if chronically short of breath)
 - *Not related to other known causes or conditions (e.g., chronic heart failure, asthma, chronic obstructive pulmonary disease)*

Other symptoms of COVID-19 can include:

- **Sore throat** (painful swallowing or difficulty swallowing)
 - *Not related to other known causes or conditions (e.g., post nasal drip, gastroesophageal reflux)*
- **Rhinorrhea** (runny nose)
 - *Not related to other known causes or conditions (e.g., returning inside from the cold, chronic sinusitis unchanged from baseline)*
- **Nasal congestion** (stuffy nose)
 - *Not related to other known causes or conditions (e.g., seasonal allergies)*

AVENUE II COVID-19 PANDEMIC PLANNING MANUAL

TOPIC: Staff Resources	SUBJECT: COVID-19 Reference Document for Symptoms Ver.	Code: SR000
		Date of Issue: September 21,2020
		Revised:
		Page: Page 2 of 3



- **New or unusual exacerbation of chronic conditions** (e.g. chronic lung diseases such as asthma, emphysema, or chronic obstructive pulmonary disorder)
- **Tachycardia** (fast heart rate), including age specific tachycardia for children
 - *Not related to other known causes or conditions (e.g., atrial fibrillation)*
- **Low blood pressure** for age
- **Hypoxia** (i.e. oxygen saturation less than 92%)
 - *Not related to other known causes or conditions (e.g., chronic obstructive pulmonary disorder)*
- **Difficulty feeding in infants**
 - *Not related to other known causes or conditions (e.g., gastroesophageal reflux disease, cleft palate)*
- **Delirium** (acutely altered mental status and inattention)
 - *Not related to other known causes or conditions (urinary tract infection, substance-related such as alcohol withdrawal, medication induced)*
- **Increased number of falls** in older persons
- **Acute functional decline** (a sudden change in ability to function compared to baseline)
 - *Not related to other known causes or conditions (e.g. gradual decline over months due to a neurological disorder such as dementia or Parkinson's disease)*

Clinical features of COVID-19 that can be diagnosed by a health care provider include:

- Clinical or radiological evidence of pneumonia

Multisystem Inflammatory Syndrome in Children and Adolescents (MIS-C) less than 19 years old

Information on this syndrome and its temporal association with COVID-19 is still emerging and may evolve over time. An assessment for MIS-C should be done by a Health Care Provider. Please see the [World Health Organization \(WHO\) Case Definition](#) or the [Canadian Paediatric Surveillance Program \(CPSP\) Case Definition](#) for diagnostic criteria.

The [WHO MIS-C preliminary case definition](#):

- Persistent fever for 3 or more days

AND two or more of the following:

AVENUE II COVID-19 PANDEMIC PLANNING MANUAL

TOPIC: Staff Resources	SUBJECT: COVID-19 Reference Document for Symptoms Ver.	Code: SR000
		Date of Issue: September 21,2020
		Revised:
		Page: Page 3 of 3



- Rash or bilateral non-purulent conjunctivitis or muco-cutaneous inflammation signs (oral, hands, or feet).
- Hypotension or shock.
- Features of myocardial dysfunction, pericarditis, valvulitis, or coronary abnormalities (including ECHO findings or elevated Troponin/NT-proBNP),
- Evidence of coagulopathy (by prolonged PT, PTT, elevated d-Dimers).
- Acute gastrointestinal symptoms (diarrhea, vomiting, or abdominal pain).

AND

- Elevated markers of inflammation such as ESR, C-reactive protein, or procalcitonin.

AND

- No other obvious microbial cause of inflammation, including bacterial sepsis, staphylococcal or streptococcal shock syndromes.

AND

- Evidence of COVID-19 (RT-PCR, antigen test or serology positive), or likely contact with patients with COVID-19